

A study on the Effects of Convective Momentum Transport Associated with Rain Bands within the Madden-Julian Oscillation

Tomoki Miyakawa ¹, Mitchell W. Moncrieff ², Tomoe Nasuno ³, Hiroaki Miura ^{1,3}, Masaki Satoh ^{1,3}, Yukari N. Takayabu ^{1,2}

1. University of Tokyo, 2. NCAR, 3. JAMSTEC


Upscale effects from unresolved Mesoscale Convective Systems (MCSs) are known sources of uncertainty in General Circulation Models (GCMs), which show general difficulty in reproduction of MJOs. The Non-hydrological Icosahedral Atmospheric Model (NICAM) successfully reproduced an MJO case using extremely fine mesh, directly resolving MCSs without cumulus parameterization.

We analyzed the upscale effect of Convective Momentum Transport (CMT) associated with rainbands of MCSs embedded within the convectively active region of the reproduced MJO case. The upscale


zonal acceleration ensemble of CMT formed a three-layered structure: positive near the surface (below 1.6km); negative at low to mid levels (2km – 6.5km); positive at upper levels (above 11km). CMT accounted for -16 m/s of the 2 km – 6.5 km averaged difference (10 m/s) of the zonal wind longitudinally across the MJO.

Two possible roles of the CMT are proposed. One is that the CMT acts to prevent eastward advection of moisture rich air, thereby delays the MJO propagation. Another is that it tilts the structure of the zonal winds associated with the MJO.


1. NICAM MJO Experiment



Icosahedral grid used in NICAM (Satoh et al. 2008)



MJO cloud clusters observed from satellite and reproduced by NICAM (Miura et al. 2007).



Composed zonal wind structure of the MJO. NICAM (above) and JGDAS-reanalysis (below). The center of the MJO is determined following the EOF based procedure described in Wheeler and Hendon (2004).


Horizontal grid	Icosahedral grid 7km (3.5km 14km)
Vertical grid	σ-coordinate, Lorenz grid 40 layers 0m ~ 38km
Duration	32 days
Governing equations	Full compressible, Nonhydrostatic
Cumulus parameterization	None
Turbulence / Surface flux	Mellor-Yamada - level 2 + water vapor
Radiation / Aerosols	MSTRNX (Sekiguchi 2004)
Cloud microphysics	Grabowski (1998)
Surface process	Bucket land model / Fixed SST
Initial condition	NCEP - Reanalysis 2006-12-15 00:00:00
Boundary conditions	Reynolds SST, Sea Ice (weekly) E-topography, Matthews vegetation UGAMP, ozone climatology (for AMP2)

Model configuration for the MJO experiment.

NICAM succeed to reproduce an MJO case observed during Dec 2006 – Jan 2007 with surprisingly high accuracy by explicitly representing mesoscale convections without cumulus parameterization (Miura et al. 2007).

In this study we use the 7km mesh output data. Three (two) dimensional data are 6 (1.5) -hourly.


2. Upscale Convective Momentum Transport within the MJO



CMT associated with rainbands embedded in MJOs cause upscale acceleration. This effect is usually parameterized as sub-grid mixing components in GCMs. However, it is known from observation and numerical experiments that organized rainbands can have counter gradient effects on the vertical wind shear (e.g. LeMone and Moncrieff, 1994; Tung and Yanai 2002).

3. Rainbands Detected in the NICAM MJO case

Threshold used for rainband detection : (Size of continuous area where precipitation ≥ 0.3mm/h) ≥ 2000km²




Rainband frequency distribution map derived from TMI and NICAM (above). The expected ratio of rainband case numbers estimated from the data retrieval difference is about 10 – 15. Red rectangle indicates the main analysis region used hereafter in this study. The lower figure shows the spatial distribution of rainbands relative to the MJO center.

An example of an ideal squall line type rainband produced in NICAM. Upper left figure shows the surface rainfall (color) and zonal wind vectors at 1580m height. Upper central figure shows the vertical wind within the region determined to be under the influence of the rainband. Upper right figure shows the vertical profile of (u, v), the horizontal wind averaged over the circle indicated in the upper left figure. The lower left figure shows vertical profiles of u (blue); v (purple); and upscale acceleration by $\frac{1}{\rho} \frac{\partial \rho u'w'}{\partial z}$ (red). The lower right figure is a zonal-height cross section of $u'w'$. The red arrow in the upper left figure indicates the zonal axis of the lower right figure.

Total number and frequency distribution of rainbands produced in NICAM are consistent with satellite data (TRMM/TMI) considering data retrieval differences. The area within 100°E - 170°E / 12°N - 12°S (red box), where the cases are abundant and the distribution agree well, is chosen as the main analysis region.


While some ideal squall line type rainbands are found (upper right figure), there were many cases with complicated structures (not shown).

4. Distribution of CMT Acceleration Vectors




MJO relative distribution of upscale CMT acceleration vectors defined by the deviation momentum flux divergence shows a three storied structure: positive zonal component near the surface (below 1.6km); negative zonal component at low to mid troposphere (2km-6.5km); positive zonal component at the upper troposphere (above 11km).

5. Quantitative Evaluation of Upscale Acceleration due to CMT



By distributing the 5° radius circles evenly over the equator, quantitative evaluation of upscale acceleration due to $\frac{1}{\rho} \frac{\partial \rho u'w'}{\partial z}$ is performed. $\frac{1}{\rho} \frac{\partial \rho u'w'}{\partial z}$ has significant contribution to the wind difference between 20° and -30° from the MJO center. In the low to mid troposphere (2km-6.5km), its contribution on the zonal wind difference longitudinally across the MJO (10 m/s) was as much as -16 m/s in average.

6. Roles of CMT ?



Miyakawa et al. 2012: Convective Momentum Transport by Rainbands within a Madden-Julian Oscillation in a Global Nonhydrostatic Model with Explicit Deep Convective Processes. Part I: Methodology and General Results, JAS