



# Improving the Representation of SGS Turbulence and Clouds in Coarse-Grid CRMs

Steve Krueger,<sup>1</sup> Pete Bogenschutz,<sup>1</sup> Marat Khairoutdinov,<sup>2</sup> Vince Larson,<sup>3</sup> and Brandon Nielsen<sup>3</sup>

<sup>1</sup>University of Utah, <sup>2</sup>State University of New York at Stony Brook, <sup>3</sup>University of Wisconsin at Milwaukee

## Multiscale Modeling Framework



## Boundary layer clouds and turbulence in deep-convection-resolving models (DCRMs)

- DCRMs are CRMs with horizontal grid sizes of 4 km or more.
- Used in MMF, GCRMs (global CRMs), and tropical cyclone models.
- In MMF and GCRMs, DCRMs are expected to represent all types of cloud systems.
- However, many cloud-scale circulations are not resolved by DCRMs.
- Representations of SGS circulations currently used in DCRMs can be improved.

### Our PDF-based parameterization

- We have constructed a 1D parameterization (Golaz et al. 2002b).
- It parameterizes layer clouds and turbulence in a unified way.
- Initially it was developed for boundary layers. It is being generalized further.
- It is based on the **Assumed PDF Method**.

### Our PDF includes several variables

We use a three-dimensional PDF of vertical velocity,  $w$ , total water (vapor + liquid) mixing ratio,  $qt$ , and liquid water potential temperature,  $\theta_l$ :

$$P = P(w, qt, \theta_l)$$

This allows us to couple subgrid interactions of vertical motions and buoyancy.

Randall et al. (1992)

### Steps in the Assumed PDF Method


The Assumed PDF Method contains 3 main steps that must be carried out for each grid box and time step:

- (1) Prognose means and various higher-order moments.
- (2) Use these moments to select a particular PDF member from the assumed functional form.
- (3) Use the selected PDF to compute many higher-order terms that need to be closed, e.g. buoyancy flux, cloud fraction, etc.

## Use results from a large-domain LES of deep convection to test the assumed PDF method

- Idealized GATE simulation with shear.
- Used SAM with 2048 x 2048 x 256 grid points and 100-m grid size for a 24-h LES.
- Equivalent to 1024 6.4-km x 6.4-km LESs.
- Collected statistics for calculating the moments needed to specify assumed PDFs for grid sizes of 800 m x 800 m x 100 m and multiples thereof.
- The statistics also include cloud fraction, liquid water mixing ratio, and its vertical flux, that can be compared to those obtained from the PDF.

## LES "visible image" 180 km x 180 km



## MISR visible image 380 km x 345 km (west of Marshall Is., 11 March 2002)








Image Credit: NASA/GSFC/LaRC/JPL, MISR Team

## PDFs of SGS cloud fraction for various horizontal grid sizes for low clouds and high clouds obtained from the large-domain LES of deep convection



## SAM & SAM-CLUBB (SAM + assumed PDF):

2D, 64-km domain, 128 levels (large-scale forcing is 6x super run)



## Summary

- We will use the "benchmark" results from a large-domain LES of deep convection to test the assumed PDF method for various horizontal grid sizes.
- We will also use the "benchmark" results to evaluate DCRMs with various configurations (SGS parameterization, grid size, domain size, and dimensionality).
- Large-domain LES of deep convection can be used to study many multiscale phenomena, such as triggering of new convection, entrainment, gust fronts, and even waterspouts.

**Acknowledgement.** This material is based upon work supported the National Science Foundation Science and Technology Center for Multi-Scale Modeling of Atmospheric Processes, managed by Colorado State University under cooperative agreement No. ATM-0425247.